The Big Thaw: gauging the past, present and future of our mountain water resources

Using novel lake-based snowfall measurements in the Rockies, Alps, and Himalayas to assess and optimise the representation of snowfall in the MetUM regional atmospheric model at kilometre grid-scales

Sid Gumber, <u>Andrew Orr</u> (anmcr@bas.ac.uk), Paul Field, Hamish Pritchard, Federico Covi, Pranab Deb, Marc Girona-Mata, Emily Potter, and Martin Widmann









Indian Institute of Technology, Kharagpur



Snow is valuable

Water and Snow Value of Water from Snow */ Puget Sound (\$) (\$) VALUE OF USABLE WATER FROM SNOW: \$11.47B ? Snowmelt here is worth up to \$90 billion per year. Current Annual Value of Usable Water from Snow \odot < \$100M \$100M to \$500M to \$1B to \$10B+ states on/off \$500M \$1B \$10B Present-day cost of water (acre/ft): \$900 \$700 \$900 \$100 \$300 \$500

Sturm et al. (2017) Water and life from snow: A trillion-dollar science question.

Snow is valuable



Snow is hazardous



Rescuers carry workers after an avalanche near Mana village in Chamoli district of India's Uttarakhand state on February 28, 2025 [Hanout/State Disaster Response Force via AFP]

Left: Aljazeera. Top right: EOS. Bottom Right: Times of India



Imja Lake-the long, silty lake in this 2010 photo-has grown in front of Imja and Lhotse Shar glaciers (top right) in the Himalayas. In 2016, engineers lowered Imja Lake by 3.5 meters to reduce the chances of a large glacial lake outburst flood. Credit: Jeffrey S. Kargel, Planetary Science Institute,

ARAKHAND Survey of 1 of '5 dangerous glacial lakes' conducted

'No Immediate Threat From Vasundhara Tal Currently, Will Be Monitored Regularly' Gauray.Talwar@timesofindia.com

Dehradun: A 15-member team comprising representatives from the Uttarakhand State Disaster Management Authority (USDMA), Indian Institute of Remote Sensing (IIRS), Wadia Institute of Himalay an Geology (WIHG), ITBP, NDRF. and SDRF conducted an assessment from Oct 15 to 22 of Vasundhara Tal a glaciall

HAZARD OF EXPANDING LAKES

> 15-member team from the Uttarakhand State Disaster Management Authority, Indian Institute of Remote Sensing, Wadia Institute of Himalayan Geology, ITBP, NDRF, and SDRF surveyed Vasundhara Tal in Chamoli from Oct 15 to 22

Assessment followed an NDMA report in Feb identifying 13 potentially hazardous glacial lakes in the state, with five, including Vasundhara Tal, marked as 'most dangerous' > Wadia Institute had said



THE TIMES OF INDIA, NEW DELHI / DEHRADUN TUESDAY, DECEMBER 3, 2024

install monitoring equipment for regular observation of the lake > Post-survey, secretary of disaster management department said there was 'no immediate threat' from the lake and

Science question: How can we constrain snowfall amount over major mountain ranges?

Science question: How can we constrain snowfall amount over major mountain ranges?

- Major data gap
- Notoriously difficult to measure (windy sites have underestimates > 50%).
- Existing in-situ measurements susceptible to bias, and most represent only a point in the landscape.
- In-situ measurements also too sparse and too poorly distributed to adequately constrain snow in weather and climate models.

Science question: How can we constrain snowfall amount over major mountain ranges?

- Major data gap
- Notoriously difficult to measure (windy sites have underestimates > 50%).
- Existing in-situ measurements susceptible to bias, and most represent only a point in the landscape.
- In-situ measurements also too sparse and too poorly distributed to adequately constrain snow in weather and climate models.

Federal Office of Meteorology and Climatology MeteoSwiss





Science question: How can we constrain snowfall amount over major mountain ranges?

- Major data gap
- Notoriously difficult to measure (windy sites have underestimates > 50%).
- Existing in-situ measurements susceptible to bias, and most represent only a point in the landscape.
- In-situ measurements also too sparse and too poorly distributed to adequately constrain snow in weather and climate models.



Precipitation correction-factors that must be applied to the observation-based "WorldClim v2" annual climatological precipitation product in order to agree with observations and models of catchment hydrology (Pritchard et al., 2021).

Current Gold Standard/ Maps of Alps and Himalayas



Accumulated total snowfall (kg/m^2) from ERA5 over the Alps (left) and west-central Himalayas (right) for the winter period of 2023-2024. Also shown are the model topographic heights.

ERA5 is the fifth generation of reanalysis data from the European Centre for Medium-Range Weather Forecasts (ECMWF), providing a comprehensive dataset of atmospheric, land surface, and ocean wave parameters. It's a global dataset that covers the period from 1940 to the present, with hourly data available. ERA5 offers a high spatial resolution of approximately 31 km and detailed vertical resolution, making it a valuable resource for climate research and monitoring

Lake-based snowfall measurements

- Measuring snowfall based on time-series of (frozen) lake water pressure (Pritchard et al., 2020)
- Lake surface area millions of times larger than pluviometer coverage



Lake-based snowfall measurements

- Measuring snowfall based on time-series of (frozen) lake water pressure (Pritchard *et al.*, 2020)
- Lake surface area millions of times larger than pluviometer coverage





Lake-based snowfall measurements





Time-series of pressure from lake-based sensor for Tomasee for the winter season from 2023-24



Time-series of pressure from lake-based sensor for Tomasee



Time-series of pressure from lake-based sensor for Tomasee



Time-series of pressure from lake-based sensor for Tomasee



Time-series of pressure from lake-based sensor and associated snowfall for Tomasee



Time-series of pressure from lake-based sensor and associated snowfall for Tomasee

Modelling snowfall using a regional atmospheric model

Model setup

- UK Met Office Unified Model (MetUM)
- Microphysics: CASIM Double Moment
- ERA5 forcing data
- Daily reinitialization approach
- Horizontal resolution: 1.5 km
- Vertical resolution: 70 levels / 40 km
- Three domains:
 - Alps (650x650 points)
 - Himalayas (650x650 points)
 - Central Rockies (USA)



Maps showing model domains and orography (m) at 1.5 km grid spacing for the (a) European Alps, (b) west-central Himalayas, and (c) central Rockies, with the positions of the lake-based snowfall measurement sites indicated by red triangles. Note that the panels have different scales.



Time series of three-hourly snowfall of observed and modelled snowfall at Tomasee (kg m^{-2}) for the period 01 Jan 2019 to 31 Mar 2019.



Control (CTRL) simulation

- Captures snowfall occurrence relatively well
- Onset/end of events well captured
- Consistent over prediction of snowfall amounts



Control (CTRL) simulation

- Captures snowfall occurrence relatively well
- Onset/end of events well captured
- Consistent over prediction of snowfall amounts

Sensitivity experiment (EXP): Reducing hydrometeor fall-speed

$$V_{x}(D) = \mathbf{a}_{x} D^{b_{x}} e^{f_{x}D} \left(\frac{\rho_{0}}{\rho}\right)^{g_{x}}$$

Where a_x , b_x , f_x and g_x represent the parameters for fall speed relationships





Sensitivity (EXP) simulation

- Reduces over-prediction / better agreement with amounts
- Onset/end of events well captured
- Precipitation sensitive to hydrometer fall-speeds



Observed Events (kg.m⁻²)

Sensitivity (EXP) simulation

- Reduces over-prediction / better agreement with amounts
- Onset/end of events well captured
- Precipitation sensitive to hydrometer fall-speeds

Scatterplots of observed versus modelled snowfall totals $(kg m^{-2})$ for all events at Tomasee. The modelled outputs are from the CTRL (black) and EXP (red) simulations. Additionally, snowfall totals from ERA5 are shown in cyan.



- Enhanced vapor deposition forming ice crystals

- Formation of graupel (in red) large ice crystals falling through supercooled cloud droplets (riming)
- Ice crystals freeze onto small snow spheres (aggregation)

Modelling snowfall / Himalayas example



Time series of three-hourly (left) and seasonal cumulative snowfall (kg m⁻²) (right) of observed and modelled snowfall at Hampta in the westcentral Himalayas for 01 Nov 2021 to 31 Mar 2022. Additionally, snowfall totals from ERA5 are shown in cyan.

Modelling snowfall / detailed maps of Alps and Himalayas



Accumulated total Snowfall (kg/m^2) from the MetUM (top) and ERA5 (bottom) for the winter period of 2023-2024. Also shown are the model topographic heights.

Employing Archer2 for high-resolution state-of-the-art snowfall modelling

SAFE User Usage Report: Dr Siddharth Gumber, (aurocumulus)

A total of 101,470 jobs were submitted by Dr Siddharth Gumber, account aurocumulus, during the period Aug 2024 - Jul 2025.



Resource	Budget	Users	Charged Usage / CUs
Archer2	n02-TBT	User 1	436,368
Archer2	n02-TBT	User 2	

SAFE User Usage Report: Dr Siddharth Gumber, (aurocumulus)



A total of 101,470 jobs were submitted by Dr Siddharth Gumber, account aurocumulus, during the period Aug 2024 - Jul 2025.

Fact: Assuming that we used all 5860 compute nodes or 750,080 cores (5860 X 2 X 64 core) continuously, that would equate to a little over **0.5815 h** of continuous full Archer2 use per user

Or using 1 full node (128 cores) continuously for about 142 days.

Cylc Workflow – Automatic Transfers from Archer to Jasmin

		command-format=rsync -aLv %(sources)s username@hpxfer4.jasmin.ac.uk:%(target)s		
Encher	- Alps	▼ F 20210122T1200Z	submitted	
		archive_files_Alps	waiting	
		archive_files_Himalayas	waiting	
		HOUSEKEEP	waiting	
More than 20 years of snowfall simulations 1999- present day	– Himalayas	🕨 🗾 glm	submitted	
		Alps_km1p5_RAL3	submitted	
		F Him_km1p5_RAL3	submitted	
		F HPC_SERIAL	waiting	
		▼ F 20210123T1200Z	running	
	Pindus	🕨 🗾 glm	submitted	
		Alps_km1p5_RAL3	running	
		Him_km1p5_RAL3	submitted	
		▼ F 20210124T1200Z	running	
		🕨 F glm	running	
		Alps_km1p5_RAL3	waiting	
		Him_km1p5_RAL3	waiting	
	SOF Disk 30TB	Limitations:		
JASMIN data facility —		 Shifting data from Disk to TAPE using ET commands and JDMA – 		
	Elastic Tape 300TB	sometimes unreliably slow – directory lockin	ig issues	
		Can keep track of requests through the ET-monitor – but opens through a		
		Jasmin machine / NoMachine client – stoppe	ed working after the new	
		Windows upgrade		
		Rogue batches from other users disturb the whole system		
	Lacks proper user interface			

Thank you

Hampta Pass Lake Indian Himalayas