


The determination of cluster structures combining infrared spectroscopy and density functional theory calculations

Piero Ferrari

Quantum Solid-State Physics

KU Leuven, Belgium


Metal clusters



A metal cluster is a particle composed of only a **countable number of metal atoms**, ranging from the dimer up to some hundreds of atoms


Size-dependent properties

Unlike bulk materials, in clusters most physical and chemical properties change drastically atom-by-atom



J. Phys. Chem. A 115, 2103 (2011)


Optical absorption



Department of Physics and Astronomy, Quantum Solid-State Physics

Size-dependent electronic structure

Clusters of specific sizes are produced in higher amounts



- Quantum confinement and electron delocalization
- Key ingredients are the geometry and the electronic structure


Molecular orbitals

KU LEUVEN


Geometry from DFT calculations

Size-dependent geometries

Bulk gold



The **size-dependency** of cluster properties is a consequence of their **geometry** and **electronic structure**




Phys. Rev. A 2018, 97, 052508

How to determine the precise geometry of an isolated cluster in a molecular beam?

The production of clusters by physical means

Laser ablation and gas condensation


"Modifications of gas aggregation sources: The double laser ablation source approach" Wiley-VCH (2017)



Department of Physics and Astronomy, Quantum Solid-State Physics


Structural information

The vibrational modes of a cluster are a fingerprint of its geometry




Infrared spectroscopy


Department of Physics and Astronomy, Quantum Solid-State Physics




Density functional theory calculations

A **structure is assigned** by comparing the experimental infrared spectrum with the calculated (harmonic) infrared spectrum of **different isomers** calculated by theory











- ✓ Calculations at PBE, Def2-TZVPP level
- Assigned structures in line with "consensus" in literature
- ✓ 2D-3D transition around n=8


The effect of doping

Pd doped Au clusters



 $\checkmark\,$ Induced 2D to 3D transition





 ✓ Magic clusters, with 6 and 8 delocalized electrons

KU LEUVEN

200

100

150


Wavenumber (cm⁻¹)




- > The technique has several advantages, as well as weak points, which must be carefully considered
- experiments, in combination with theoretical calculations


> The size-dependent properties of clusters depend on the precise geometry

> This geometry can be determined by infrared multiple photon spectroscopy



PdAu₃Ar₄







of the particle

Conclusions

Acknowledgement

KU LEUVEN

KU Leuven

Prof. Ewald Janssens Prof. Peter Lievens Dr. Jan Vanbuel





Financial Support

Research Foundation - Flanders (FWO)





Fritz-Haber-Institut der Max-Planck-Gesellschaft, Germany

MAX-PLANCK-GESELLSCHAFT

Dr. André Fielicke





Radboud University Nijmegen

Dr. Joost M. Bakker Olga Lushchikova





King's College London

Dr. Francesca Baletto Laia Delgado Callicó




Department of Physics and Astronomy, Quantum Solid-State Physics





Thanks for your attention

Department of Physics and Astronomy, Quantum Solid-State Physics

Benchmark analysis







Au₈Ar₃*

lso 1

lso 2

0.03 eV

0.00 eV

Wavenumber (cm⁻¹)

Department of Physics and Astronomy, Quantum Solid-State Physics

The effect of Ar

